

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

LUTESCENT WARBLER, Helminthophila celata lutescens. — First seen April 2, when it became common. It breeds here. I have found one nest, which was placed on the ground and contained four young The nest was composed of dry leaves and grass and lined with horse hair.

YELLOW WARBLER, *Dendroica æstiva*.—This is not a common warbler. First seen May 8. It is more domestic in its habits than the other warblers. I have not yet found a nest, though it breeds here.

WESTERN YELLOW-THROAT, Geothlypis trichas occidentalis.—This is a common warbler in low ground. First seen April 4, but did not become numerous until April 12. Habits and song like eastern bird. I have not yet found the nest.

HERMIT WARBLER, *Dendroica occidentalis*.—Is said to occur here, but I have never found it in three years of collecting.

George D. Peck, Salem, Oregon.

INFORMATION WANTED.

A monograph of the Flicker, *Colaptes auratus*, could scarcely lay any claims toward completeness without a more or less full list of the various local names or "vernacular synonyms" as they have been termed. No American bird can approach our subject in the number and variety of its nicknames. Already the number recorded is large, and, no doubt, as many more are in use locally or throughout extensive sections of this broad land of ours, of which no record has ever been made.

It is our purpose to make the list as full as possible, and in whatever language or dialect the name may be, it would be gladly received as an addition to the general knowledge of the subject.

Any information or suggestions relating to the meaning or probable origin of the following names will be gratefully received: Cave-duc (Maine), Fiddler (Cape Cod, Mass.), Harry-wicket or Hairy-wicket (New England), Hivel (New York, Ohio), Hittock or Hittucks (Canada), Ome-tuc (Maine), Talpa-na-ni, Seminole (Southern Florida), Will Crisson (South), Wood-quoi (Conn.), Yacker and Yucker (New York), Zebec (locality unknown).

Anyone who has heard of this species being designated by such names as Rainfowl, Woodpie, Woodnacker, Wild Guinea, or appellations of like nature, would confer a favor by communicating with the undersigned.

All who have promised data or desire to contribute to the Flicker Bulletin, are requested to send in their notes as early as possible.

FRANK L. BURNS, Berzeyn, Penn.